

Autism

What is Autism Spectrum Disorder?

[Español \(Spanish\)](#)

Autism spectrum disorder (ASD) is a developmental disability that can cause significant social, communication and behavioral challenges. There is often nothing about how people with ASD look that sets them apart from other people, but people with ASD may communicate, interact, behave, and learn in ways that are different from most other people. The learning, thinking, and problem-solving abilities of people with ASD can range from gifted to severely challenged. Some people with ASD need a lot of help in their daily lives; others need less.

A diagnosis of ASD now includes several conditions that used to be diagnosed separately: autistic disorder, pervasive developmental disorder not otherwise specified (PDD-NOS), and Asperger syndrome. These conditions are now all called autism spectrum disorder.

Signs and Symptoms

People with ASD often have problems with social, emotional, and communication skills. They might repeat certain behaviors and might not want change in their daily activities. Many people with ASD also have different ways of learning, paying attention, or reacting to things. Signs of ASD begin during early childhood and typically last throughout a person's life.

Children or adults with ASD might:

- not point at objects to show interest (for example, not point at an airplane flying over)
- not look at objects when another person points at them
- have trouble relating to others or not have an interest in other people at all
- avoid eye contact and want to be alone

- have trouble understanding other people's feelings or talking about their own feelings
- prefer not to be held or cuddled, or might cuddle only when they want to
- appear to be unaware when people talk to them, but respond to other sounds
- be very interested in people, but not know how to talk, play, or relate to them
- repeat or echo words or phrases said to them, or repeat words or phrases in place of normal language
- have trouble expressing their needs using typical words or motions
- not play "pretend" games (for example, not pretend to "feed" a doll)
- repeat actions over and over again
- have trouble adapting when a routine changes
- have unusual reactions to the way things smell, taste, look, feel, or sound
- lose skills they once had (for example, stop saying words they were using)

[Learn about developmental milestones that young children should reach »](#)

Diagnosis

Diagnosing ASD can be difficult since there is no medical test, like a blood test, to diagnose the disorders. Doctors look at the child's behavior and development to make a diagnosis.

ASD can sometimes be detected at 18 months or younger. By age 2, a diagnosis by an experienced professional can be considered very reliable.¹ However, many children do not receive a final diagnosis until much older. This delay means that children with ASD might not get the early help they need.

Treatment

There is currently no cure for ASD. However, research shows that early intervention treatment services can improve a child's development. Early intervention services help children from birth to 3 years old (36 months) learn important skills. Services can include therapy to help the child talk, walk, and interact with others. Therefore, it is important to talk to your child's doctor as soon as possible if you think your child has ASD or other developmental problem.

Even if your child has not been diagnosed with an ASD, he or she may be eligible for early intervention treatment services. The [Individuals with Disabilities Education Act \(IDEA\)](#) says that children under the age of 3 years (36 months) who are at risk of having developmental delays may be eligible for services. These services are provided through an early intervention system in your state. Through this system, you can ask for an evaluation.

In addition, treatment for particular symptoms, such as speech therapy for language delays, often does not need to wait for a formal ASD diagnosis.

Causes and Risk Factors

We do not know all of the causes of ASD. However, we have learned that there are likely many causes for multiple types of ASD. There may be many different factors that make a child more likely to have an ASD, including environmental, biologic and genetic factors.

- Most scientists agree that genes are one of the risk factors that can make a person more likely to develop ASD.
- Children who have a sibling with ASD are at a higher risk of also having ASD.
- Individuals with certain genetic or chromosomal conditions, such as fragile X syndrome or tuberous sclerosis, can have a greater chance of having ASD.
- When taken during pregnancy, the prescription drugs valproic acid and thalidomide have been linked with a higher risk of ASD.
- There is some evidence that the critical period for developing ASD occurs before, during, and immediately after birth.
- Children born to older parents are at greater risk for having ASD.

ASD continues to be an important public health concern. Like the many families living with ASD, CDC wants to find out what causes the disorder. Understanding the factors that make a person more likely to develop ASD will help us learn more about the causes. We are currently working on one of the largest U.S. studies to date, called [Study to Explore Early Development \(SEED\)](#). SEED is looking at many possible risk factors for ASD, including genetic, environmental, pregnancy, and behavioral factors.

Who is Affected

ASD occurs in all racial, ethnic, and socioeconomic groups, but is about 4 times more common among boys than among girls.

For over a decade, CDC's [Autism and Developmental Disabilities Monitoring \(ADDM\) Network](#) has been estimating the number of children with ASD in the United States. We have learned a lot about how many U. S. children have ASD. It will be important to use the same methods to track how the number of children with ASD is changing over time in order to learn more about the disorder.

If You're Concerned

If you think your child might have ASD or you think there could be a problem with the way your child plays, learns, speaks, or acts, **contact your child's doctor, and share your concerns.**

If you or the doctor is still concerned, **ask the doctor for a referral to a specialist** who can do a more in-depth evaluation of your child. Specialists who can do a more in-depth evaluation and make a diagnosis include:

- Developmental Pediatricians (doctors who have special training in child development and children with special needs)
- Child Neurologists (doctors who work on the brain, spine, and nerves)
- Child Psychologists or Psychiatrists (doctors who know about the human mind)

At the same time, call your state's public early childhood system to request a free evaluation to find out if your child qualifies for intervention services. This is sometimes called a Child Find evaluation. You do not need to wait for a doctor's referral or a medical diagnosis to make this call.

Where to call for a free evaluation from the state depends on your child's age:

- **If your child is not yet 3 years old, contact your local early intervention system.**
 - You can find the right contact information for your state by calling the Early Childhood Technical Assistance Center (ECTA) at 919-962-2001.
 - Or visit the [ECTA website](#).

- **If your child is 3 years old or older, contact your local public school system.**
 - Even if your child is not yet old enough for kindergarten or enrolled in a public school, call your local elementary school or board of education and ask to speak with someone who can help you have your child evaluated.
 - If you're not sure who to contact, call the Early Childhood Technical Assistance Center (ECTA) at 919-962-2001.
 - Or visit the [ECTA website](#).

Research shows that early intervention services can greatly improve a child's development.^{2,3} In order to make sure your child reaches his or her full potential, it is very important to get help for an ASD as soon as possible.

Vaccine Safety

Some people have had concerns that ASD might be linked to the vaccines children receive, but studies have shown that there is no link between receiving vaccines and developing ASD. For more information about vaccines and ASD, [click here](#)

For more information go to the [CDC Autism Spectrum Disorder page](#).

Article Source

Centers for Disease Control (CDC)

Source URL

<https://www.cdc.gov>

Last Reviewed

Saturday, October 17, 2020